

Cal Poly Humboldt

## Digital Commons @ Cal Poly Humboldt

---

IdeaFest 2023

---

2023

### Effects of Tidal Conditions on Egret Foraging Behavior in Arcata Marsh

Tania Estrada Rodriguez

*Cal Poly Humboldt*, tlt11@humboldt.edu

Follow this and additional works at: <https://digitalcommons.humboldt.edu/ideafest2023>

---

#### Recommended Citation

Estrada Rodriguez, Tania, "Effects of Tidal Conditions on Egret Foraging Behavior in Arcata Marsh" (2023). *IdeaFest 2023*. 98.

<https://digitalcommons.humboldt.edu/ideafest2023/98>

This Article is brought to you for free and open access by Digital Commons @ Cal Poly Humboldt. It has been accepted for inclusion in IdeaFest 2023 by an authorized administrator of Digital Commons @ Cal Poly Humboldt. For more information, please contact [kyle.morgan@humboldt.edu](mailto:kyle.morgan@humboldt.edu).

# Tidal Nonsense

## Effects of Tidal Conditions on Egret Foraging Behavior in Arcata Marsh

Tania L. Estrada Rodriguez, California Polytechnic State University, Humboldt, 1 Harpst Street Arcata Ca, 95521, USA

### Introduction

- Egrets regulate wetland ecosystems by regulating populations of fish and invertebrates, but the impact of tide levels on their foraging success is poorly understood.
- Tide levels affect egret populations, and environmental conditions influence their foraging behavior (Strong et al. 1997, Maccrone and Brzorad 2002).
- **Objective:** to determine if high tides pose unfavorable foraging conditions for egrets and if their foraging behavior differs between high and low tide

### Hypothesis


- If high tides pose unfavorable foraging conditions for egrets, such as by reducing the visibility of prey, then we **expect** to observe greater foraging behaviors like strike rates during low tide

### Methods

- **Sample size:** 6 ponds labeled: 1 - Brackish Pond, 2 - Gearheart Marsh, 3 - Hauser Marsh, 4 - Allen Marsh, 5 - Restored Salt Marsh, and 6 - Log Pond.
- **Variables:** tide levels, temperature, precipitation, strike rates, time spent foraging.
- **Observations:** 10-minute focal observations to minimize observer bias or interference with the bird's natural behavior.
- **Analysis:** T-test and Logistic regression in R statistical software

**Outcome variables:** Strike rate

**Predictor variable:** Temperature, Overcast, Waterfowl presence, Tide



### Results

- tide did not impact egret foraging behavior  
( $T = 1.730$ ,  $DF = 95\%$ ,  $P = 0.086$ ).
- Logistic regression analysis compared AIC values, mod2 was lower at 129.09 than mod1 at 130.91
- my top-performing model showed inverse relationship between weather and egret foraging success, with Beta coefficient of -1.10 and a significant p-value of 0.03.

### Discussion

- Contradicts previous studies (Lantz et al. 2010) Low tides were preferred
- Mean proportion of foraging time was similar for both groups.
- Weather extremes reduced egret activity at the AMWS: hail, snow, wind, rain (Youcefi et al. 2018)
- Future research could examine individual species and locations.

### Literature cited

- Strong A. M., G. T. Bancroft, and S. D. Jewell. 1997. Hydrological constraints on tricolored heron and snowy egret resource use. *Condor* 99: 894-905.
- Maccarone A.D., and J.N. Brzorad. 2002. Foraging Patterns of Breeding Egrets at Coastal and Interior Locations. *Waterbirds*, 25.
- Martin, P., and P. Bateson. 2007. *Measuring Behaviour: an introductory guide*. Cambridge University Press.
- Lantz, S. M., Gawlik, D. E., & Cook, M. I. (2010). The Effects of Water Depth and Emergent Vegetation on Foraging Success and Habitat Selection of Wading Birds in the Everglades. *Wetlands*, 30(6), 1163-1173. doi: 10.1007/s13157-010-0124-7.
- Youcefi, A., Baaloudj, A., Zebba, R., Mahdjoub, H., Bensakhri, Z., & Bensouilah, S. (2018). Weather conditions affect the collective roosting behaviour of the Cattle Egret *Bubulcus ibis*. *Avian Research*, 9(1), 9. doi:10.1186/s40657-018-0091-6.