

Cal Poly Humboldt

Digital Commons @ Cal Poly Humboldt

Botanical Studies

Open Educational Resources and Data

2022

Purposeful Use Of Poisonous Plants

James P. Smith Jr

Follow this and additional works at: https://digitalcommons.humboldt.edu/botany_jps



Part of the [Botany Commons](#)

PURPOSEFUL USE OF POISONOUS PLANTS

James P. Smith, Jr.
Professor Emeritus of Botany
Department of Biological Sciences
Humboldt State University
Arcata, California

11 April 2021

Words like “poisonous” and “toxic” tend to conjure up negative images of illness, pain, and death. In humans the encounters with poisonous plants often involve mistaken identity or lack of knowledge about the plant’s capability. But a more comprehensive look at our interaction with poisonous plants must take into account that we have knowingly used them as instruments of punishment, torture, murder, and suicide. We have also used our knowledge of plant toxins to make arrow poisons and to kill fish, insects, and rodents. Probably the least known and most mysterious is our use of these plants by various societies in ordeal rituals that determine guilt. All of these are the subject of this paper.

On the much more positive side of the ledger, many of our best known medicines come from plants where we take advantage of their toxic effects to achieve the desired effects in controlled situations. Many people are alive today because of the glycosides in the foxglove plant and their effect on the contraction of the heart.

ARROW & DART POISONS

"We did not escape without damage because [the Indians] killed another companion of ours ... and in truth, the arrow did not penetrate half a finger, but as it had poison on it, he gave up his soul to our Lord." (Francisco de Orellana, 1541)

Peoples of both the Old World and the New World have used many different plants in the preparation of these poisons. Native Americans on this continent used relatively few arrow poisons from plant sources, relying instead on rattlesnake venom and the juices of the black widow spider. The Indians of South America and the tribesmen of Africa used arrow poisons both to hunt wild game and in wars against their enemies.

CURARE. This is probably the most famous arrow poison. The name is a phonetic rendering of an Indian phrase meaning, "He, to whom it comes, falls." Other spellings used in the literature are urari, woorari, woorali, and wourali. Curare is the name for a whole group of arrow poisons used in South America. The fact that the ingredients and their relative proportions vary from one location to the next hindered any real understanding of the botanical and chemical nature of the poison.

The Baron Alexander von Humboldt was probably the first European to witness the preparation of curare. He wrote that the Indians shredded bark of certain trees and made an infusion from it. This was further concentrated by boiling it. Several other plant materials were added to the mixture.

The Indians have several different ways of assaying the strength of curare. One is to wound a frog with a poisoned arrow or dart. If it can jump more than eight times without the curare taking effect, then the potion is deemed too weak. Another test involves a monkey jumping from tree to tree. When the curare is at its proper strength, an animal should not be able to jump to more than one tree before the toxin takes effect. And, a final test is that since the honor of preparing curare often falls to the old women of the tribe, if they are not half intoxicated by the fumes of the boiling mixture, the brew is not yet ready.

The exact botanical nature of curare is still in some doubt. The two principal toxic ingredients are *Strychnos* spp. (often *S. toxifera*) of *Loganiaceae* and *Chondrodendron tomentosum* of the moonseed family (*Menispermaceae*).

Three basic groups or kinds of curare are often recognized, based upon the physical form in which they are prepared. **Tubocurare**, also known as tube curare or bamboo curare, is cylindrical, having been packed in hollow bamboo stems. The two other kinds are **gourd** or calabash curare and **pot** curare. *Chondrodendron* is the principal ingredient in tube curare; it is also used in pot curare. *Strychnos* is used in both calabash and pot curare.

The active ingredient is d-tubocurarine chloride or tubocurarine chloride, a whitish, odorless powder. Poisoning manifests itself in muscular relaxation by blocking of impulses between the nerve and the muscle fiber activated by it. Symptoms include impaired vision; bilateral drooping of lips; heaviness of face; relaxation of

jaw; weakness of head muscles; inability to raise head; paralysis of spinal muscles, legs, & arms; slowed respiration; and death. Curare is rapidly destroyed and excreted. Its action usually lasts for about 15-20 minutes, during which the victim dies. The flesh of animals killed by curare may be eaten.

There are also medicinal uses for curare. Because it is a muscle relaxant, it may be used in the treatment of convulsive mental patients and in certain types of surgery. Curare is also used to diagnose myasthenia gravis, a muscular disorder characterized by an overall deterioration of muscle tone.

KOMBI. The action of this African arrow poison was first described in detail by Dr. David Livingstone, the missionary to Africa of "Dr. Livingstone, I presume" fame. He observed people preparing a poison from the fruits of *Strophanthus*, a member of Apocynaceae, the dogbane family. The action of kombi is much like that of curare. The wounded animal rarely runs more than about a hundred yards before it falls. Kombi contains strophanthin, a glycoside with digitalis-like effects. It must get into the blood stream, which means that the flesh of kombi victims is edible.

Strophanthin also finds limited uses in medicine. It is used on heart attack victims when time is of the utmost importance. It takes effect much faster than digitalis, but it is such a violent poison that the dosages must be carefully controlled.

UPAS. This is an arrow poison used by the natives of Java. The chief ingredient is *Antiaris toxicaria* of Moraceae, the mulberry family.

OUABAIN. The natives of tropical Africa prepare an arrow poison from *Strophanthus gratus*. A somewhat inferior brand can be made from *Acokanthera schimperi* and *A. deflexa*. Ouabain is the most rapidly acting heart-glycoside known.

NEW WORLD ARROW & DART POISONS

Scientific Name (Family)	Area of Use
<i>Aconitum</i> spp. • Buttercup	North America
<i>Anemone</i> spp. • Buttercup	North America
<i>Anomospermum</i> spp. • Moonseed	South America
<i>Capsicum annuum</i> • Nightshade	South America
<i>Cocculus</i> spp. • Moonseed	South America
<i>Colliguaja odorifera</i> • Spurge	South America
<i>Chondrodendron</i> spp. • Moonseed	South America
<i>Conium maculatum</i> • Parsley	North America
<i>Cynanchum macrophyllum</i> • Milkweed	North America
<i>Delphinium elatum</i> • Buttercup	North America
<i>Dieffenbachia seguine</i> • Philodendron	Cuba
<i>Dioscorea</i> spp. • Yam	West Indies
<i>Ficus atrox</i> • Mulberry	South America
<i>Gutteria venificiorum</i> • Annona	South America
<i>Hippomane mancinella</i> • Spurge	West Indies
<i>Hura crepitans</i> • Spurge	Central & South America
<i>Jacquinia</i> spp. • Theophrasta	Central & South America
<i>Nicotiana</i> spp. • Nightshade	Central & South America
<i>Papaver nudicaule</i> • Poppy	North America
<i>Pedilanthus tithymaloides</i> • Spurge	West Indies
<i>Piper geniculatum</i> • Pepper	South America
<i>Piscidia piscipula</i> • Bean	North America
<i>Ranunculus</i> spp. • Buttercup	North America
<i>Sapium biloculare</i> • Spurge	Mexico
<i>Sebastiania palmeri</i> • Spurge	Mexico
<i>Serjania</i> spp. • Soapberry	West Indies
<i>Spigelia fruticulosa</i> • Logania	South America
<i>Strychnos</i> spp. • Logania	South America
<i>Tabernaemontana</i> spp. • Dogbane	South America
<i>Yucca glauca</i> • Lily	North America

FISH POISONS

Many different plants have been used to poison, or more precisely, to stupefy fish. The technique is reasonable straightforward. It typically involves putting the required part of a particular plant in a still body of water. The plant toxin is released into the water, numbing the fish so that it comes floating to the surface, where they can be gathered up.

The California buckeye (*Aesculus californica*) was used by several tribes of Native Americans. Seeds,

sometimes leaves or young stems, were crushed and made into a mash that was floated on the surface of the water. The stupefied fish were then gathered by hand or in nets.

The fish poison tree (*Piscidia piscipula*) of the bean family occurs in Texas, Florida, Mexico, the West Indies, and South America. Its leaves, stems, and root bark are macerated with a rum distillation residue or lime water. The material is then placed in baskets and floated in the water until the stupefied fish surface. The plant contains piscidin, a mixture of two toxic glycosides.

Scientific Name • Common Name	Where Used
<i>Aesculus californica</i> • California buckeye	California
<i>Chlorogalum pomeridianum</i> • soaproot	California
<i>Clibadium</i> spp. • Cunambi	Amazon
<i>Conium maculatum</i> • wasia	Mexico
<i>Croton setigerus</i> • turkey mullein	California
<i>Derris elliptica</i> • tuba-root	Asia
<i>Echinocystis fabacea</i> • manroot	California
<i>Erythrina piscipula</i>	South America
<i>Hura crepitans</i> • oassucú	South America
<i>Lomatium dissectum</i>	California
<i>Lonchocarpus</i> spp. • barbasco	Tropical America
<i>Lycopus</i> spp. • horehound	California
<i>Manihot esculenta</i> • cassava	Guiana
<i>Patinoa sphaerocarpa</i>	Amazon
<i>Paullinia</i> spp. • timbó	South America
<i>Phyllanthus</i> spp.	South America
<i>Sapindus saponarius</i> • amolli	Mexico
<i>Serjania</i> spp. • timbó	South America
<i>Smilacina sessilifolia</i> • Solomon seal	California
<i>Tephrosia piscatoria</i>	South America
<i>Thevetia peruviana</i> • jorro-jorro	Brazil
<i>Trichstema lanceolatum</i> • blue curls	California
<i>Umbellularia californica</i> • pepperwood	California
<i>Wikstroemia</i> spp.	Hawai'i

INSECTICIDES

In addition to the more exotic arrow and ordeal poisons, the plant kingdom yields useful poisons for the control of insects. About 1200 different species have been used in this context, although only rotenone, pyrethrum, and nicotine are of any real commercial value. One of these, rotenone, was used for centuries as a fish poison in South America before its use as an insecticide.

Rotenone is a very popular insecticide derived from the roots of *Derris* and *Lonchocarpus*, both members of the bean family. The former is native to the Old World tropics, while the latter is New World in its distribution. Rotenone was first isolated from *Derris* in 1902. It is a colorless, crystalline compound extracted from the dried roots. Rotenone is insoluble in water, but soluble in oil. It is about fifteen times more toxic than nicotine. The toxic principle has no effect on warm blooded animals. Rotenone was isolated from the roots of *Lonchocarpus* in 1926. Its roots are richer in the toxin than those of *Derris*.

Pyrethrum is the name given to the dried flower heads of three species of *Chrysanthemum*. They yield volatile oils that paralyze insects. This came as no surprise to horticulturists who knew that these plants were rarely attacked by insects. Depending upon the species, the pyrethrum is often called "Dalmatian insect powder," "Caucasian insect powder," or "Persian insect powder." Pyrethrum has been used successfully in the control of flies, fleas, malarial mosquitos, and body lice.

Scientific Name • Common Name	Comment
<i>Anabasis ioaphylla</i>	Used in Near East
<i>Annona cherimola</i> • cherimoya	Used against lice
<i>Chrysanthemum cinerariifolium</i>	Pyrethrum
<i>Croton texensis</i> • croton	Used in New Mexico
<i>Cymbopogon nardus</i> • lemon grass	Citronella oil
<i>Delphinium</i> spp. • larkspur	Seeds used
<i>Derris</i> spp. • tuba-root	Root source of rotenone
<i>Dolichos pseudopachyrrhizus</i> • hyacinth bean	Tropical Africa
<i>Duboisia hopwoodii</i>	Leaves contain nornicotine
<i>Eupatorium</i> spp. • boneset	Eastern N. America
<i>Gardenia lucida</i>	Used in Asia to ward off flies
<i>Ipomoea quamoclit</i> • morning glory	Seeds used
<i>Liquidambar styraciflua</i> • sweet gum	Fumigant
<i>Lonchocarpus</i> spp. • barbasco	Source of rotenone

<i>Macuna</i> spp. • velvet bean	Seeds contain L-dopa
<i>Melia azedarach</i> • China berry	Used against flies
<i>Nicotiana</i> spp. • tobacco	Leaf dustings used
<i>Pinus palustris</i> • long-leaved pine	Pine oil used
<i>Ruta graveolens</i> • rue	Leaves contain volatile oil
<i>Schoenocaulon officinale</i> • sabadilla	Seeds used
<i>Sesamum indicum</i> • sesame	Used with pyrethrum
<i>Tanacetum vulgare</i> • tansy	Used on fleas and lice
<i>Xanthoxylum clava-herculis</i> • prickly-ash	Bark

RODENTICIDES

These plant toxins are used in poison baits, primarily to kill rats. Therefore, they are sometimes called **raticides**. They are effective in small quantities, and appear to taste just fine.

Squill or sea-onion • *Drimia maritima* is a Mediterranean member of the lily family. Since ancient times, its bulbs and extracts of the bulbs have been used to kill rodents. If you or I eat this material, it will make us sick and we will vomit, thereby ridding the system of the toxin. Rodents, on the other hand, have no “vomit control center” in their brains so that any plant parts consumed will remain in the stomach and be absorbed. The sea-onion contains a series of toxins that affect the heart. They are called cardiac glycosides and they are chemically similar to those found in the foxglove plants. The sea-onion toxins, given by injection in moderation, have medicinal uses as diuretics, emetics, and expectorants.

Sweet clover. One of the most pleasant smells of rural areas is freshly mown pastures. Part of that nice aroma comes from sweet clovers • *Mellilotus* spp., herbs of the legume family. These plants also produce a form of **coumarin**, which inhibits blood clotting. Given in controlled quantities, coumarins are effective as blood thinners. Some years ago, the Wisconsin Alumni Research Foundation discovered that a modified form of coumarin made a very effective rat poison. They named it **warfarin** – after themselves. The poor little rats got off someplace and die from internal hemorrhaging.

Scientific Name (Common Name)	Comment
<i>Coriaria thymifolia</i> (shanshi)	Americas & Pacific Islands; psychoactive
<i>Dieffenbachia seguine</i> (dumb cane)	Also a popular, but toxic ornamental
<i>Drimia maritima</i> (red squill, sea-onion)	native to Old World
<i>Gliricidia sepium</i> (quick stick)	Shade crop for coffee; also insecticide
<i>Helleborus niger</i> (black-hellebore, Christmas-rose)	Also a powerful purgative
<i>Persea americana</i> (avocado, alligator-pear)	Also toxic to house pets
<i>Spondianthus preussii</i>	Also a fish poison and in executions
<i>Veratrum album</i> (white-hellebore)	Native to Europe & Asia

ORDEAL POISONS

The lives of many peoples through the ages have been dominated by a complex pantheon of deities, both good and evil. In many societies, one very important obligation was to placate friendly gods and not to encourage the evil ones. These were not societies that were protected by a single omnipotent god who rewarded good deeds and punished the wicked. In many tribes, the people developed procedures for determining guilt and innocence. This was a trial or an ordeal to which suspects were subjected. If innocent, the person would be able to pass the ordeal; but, if guilty, he or she would fail.

We have used similar devices in the United States and Europe. Consider the witch trials of the 17th century. A woman suspected of being a witch was bound, placed in a bag, and tossed in a lake. If she floated, it was concluded that she was indeed a witch and she was dispatched in some novel fashion. If the woman sank to the bottom of the lake and drowned, her survivors had the consolation of knowing that she was innocent.

Many of the African tribes employed plant poisons in their own version of the ordeal. No one really knows how the use of ordeal poisons began. One plausible explanation is that food gatherers mistakenly picked a toxic plant, assuming it to be edible. The people of the village ate it; some were poisoned and died, while others lived. Given the philosophical and religious climate, a possible explanation that would come to mind is that those who ate the plant and died were somehow connected with witchcraft or were guilty of some evil and were being punished.

There are several different ways in which the ordeal poisons were employed. One of the most straightforward involves a plant called the **tanghin of Madagascar** • *Cerbera* spp. All parts of this tree are toxic, the fruits especially so. The toxin is a cardiac glycoside. The person undergoing the ordeal is given a drink of rice water or rice soup. He or she is then given three pieces of chicken skin to swallow without chewing, followed by the crushed tanghin fruits mixed with banana juice and a preparation of cardamom leaves or juice. Incantations are performed. The concoction acts as an emetic, a substance that will cause vomiting. Failure to do so is an immediate sign of guilt. The vomitus is carefully examined by tribal officials to see if all three pieces of chicken

skin are present. They must be if the suspect is to be judged innocent. If not, the other members of the tribe kill the person, knowing now that he or she is guilty. The significance of the chicken skin lies in the belief that an evil spirit inhabits the body of a guilty person and that it would devour any flesh presented to it, including chicken skin. Before considering this to be a quaint custom, consider that in 1830 a series of natural disasters and diseases swept Madagascar. This prompted mass ordeal trials to determine who was causing these calamities. As many as 6000 people died in one trial!

Probably the most famous of the ordeal poisons involves the **Calabar bean** • *Physostigma venenosum*, a vine of the legume family named after a town in Nigeria. It has a long history of use in Africa as a means of identifying and killing witches. The trial consisted of the accused drinking water that contained eight smashed Calabar beans. If the person regurgitated the beans or raised his right arm and survived, then he was judged innocent. The fast-acting toxin, **physostigmine**, rarely produced that result.

In recent years this plant has been the object of medical research. This same toxin is now used to treat glaucoma, chronic constipation, and aspects of Alzheimer's disease.

The "**Mauvi tree**" of East Africa is used by the natives of the Lake Nyassa region. The bark is mixed with other substances and given to the suspects. If you are guilty, you will vomit and die. There is a certain simplicity to it. The body of the guilty party is spirited away by relatives who will cremate the remains, grind up the bones, and put the powder on their faces.

In some instances, the suspects at an ordeal will be asked to perform some task. Inability to do so is taken as a sign of guilt. In one tribe, the medicine man draws a line on the ground about ten feet in front of the suspects. If you can walk across the line, you are innocent. In another tribe, the suspect is asked to jump over a stick held about 50 cm above the ground. One tribe that occupied what used to be called the Ivory Coast in Africa used the white latex from a member of the spurge family. A latex-soaked piece of cotton was placed under the eyelid and allowed to remain there for a required period. Damage to the cornea was taken as evidence of guilt.

Many of the ordeal poisons were simple preparations involving a toxic plant and one or two other ingredients. Some were complex preparation of a plant poison, lizards, crushed snakes, toads, hearts from previous victims, miscellaneous human organs, blood, and bile. The various ingredients were then fermented for one year to bring them up to proper strength.

The belief in the accuracy of the ordeal poisons was so powerful that perfectly innocent people who were subjected to the ritual would confess to the most horrible crimes if they felt the toxin taking effect. As with so many useful social institutions, corruption soon tainted the ordeal rituals. The priests and witch doctors found that they could guarantee the outcome of an ordeal by careful preparation of the poisons. Their friends received weak doses; their enemies seldom survived. The witch doctor would make a conspicuous display of drinking a weak dose of the ordeal poison before a large crowd to strengthen his position in the community and to show that the innocent had nothing to fear.

PLANTS USED IN ORDEAL RITUALS

Common/scientific Name	Where Used?
African pearwood • <i>Mimusops djave</i>	Africa
Bushman's poison • <i>Akokanthera venetata</i>	Africa
Calabar bean • <i>Physostigma venenosum</i>	Africa
<i>Combretum confertum</i>	Africa
Desert-rose • <i>Adenium obesum</i>	Africa/Arabia
Sassy bark • <i>Erythrophleum suaveolens</i>	Old World tropics
<i>Strychnos icaia</i>	Africa
<i>Strychnos kipapa</i>	Africa
Tallow tree • <i>Detarium senegalense</i>	Africa
Tanghin of Madagascar • <i>Cerbera vinenifera</i>	Indian/Pacific coasts
Violet-tree • <i>Securidaca longipedunculata</i>	Tropical Africa

PLANTS USED IN EXECUTIONS

Common/scientific Name	Where Used?
Allspice jasmine • <i>Gelsemium elegans</i>	Indomalaysia
Bejuco de agua • <i>Rourea glabra</i>	C. & S. America
Hemlock water-dropwort • <i>Oenanthe crocata</i>	Europe
Mexican shrubby-spurge • <i>Euphorbia cotinifolia</i>	C. & S. America
Monk's hood • <i>Aconitum napellus</i>	Europe
Muavi tree • <i>Parkia bussei</i>	Africa
Oleander • <i>Nerium oleander</i>	Mediterranean
Pink root • <i>Spigelia anthelmia</i>	Old World tropics
Poison hemlock • <i>Conium maculatum</i>	Greece

Rat's bane • *Dichapetalum toxicaria*
 Sago palm • *Metroxylon sagu*
 Timbo • *Paullinia pinnata*
 Yellow heads • *Gnidia krausiana*

Africa
 Malaysia
 Mexico & C. America
 Africa

PLANTS USED TO COMMIT SUICIDE

Common/scientific Name

Kaffir-onion • *Boöphane disticha*
 Tanghin of Madagascar • *Cerbera manghas*
 Allspice jasmine • *Gelsemium elegans*
 Glory-lily • *Gloriosa superba*
 Jequirity or rosary bean • *Abrus precatorius*
 Mountain-laurel • *Kalmia latifolia*
Phyllanthus engleri
 Catchbird tree • *Pisonia brunoniana*

Where Used?

Africa
 Indian/Pacific coasts
 Indomalaysia
 India
 E. North America
 Africa
 Tropical Asia/Oceania

BIBLIOGRAPHY

ARROW & DART POISONS

Bisset, N. G. 1966. The arrow and dart poisons of South-East Asia, with particular reference to the *Strychnos* species used in them. Part I. Indonesia, Borneo, Philippines, Hainan, and Indo-China. *Lloydia* 29: 1-13.

Bisset, N. G. 1981. Arrow poisons in China. II. *Aconitum* -- botany, chemistry, and pharmacology. *J. Ethnopharm.* 4: 247-336.

Bisset, N. G. 1989. Arrow and dart poisons. *J. Ethnopharm.* 25: 1-41.

Bisset, N. G. 1992. War and hunting poisons in the New World. Pt. 1. Notes on the early history of curare. *J. Ethnopharm.* 36(1): 1-26.

Bisset, N. G. & A. J. M. Leeuwenberg. 1968. The use of *Strychnos* species in Central African ordeal and arrow poisons. *Lloydia* 31: 208.

Bisset, N. G. & M. C. Woods. 1966. The arrow and dart poisons of South-East Asia, with particular reference to the *Strychnos* species used in them. II. Burma, Thailand and Malaya. *Lloydia* 29: 172-.

Blubaugh, L. V. & C. R. Lineger. 1948. Curare and modern medicine. *Econ. Bot.* 2: 73-82.

Bovet, D. F. Bovet-Nitti, & G. B. Marini-Bettolo. 1959. Curare and curare-like agents. Van Nostrand. Princeton, MA.

Bradley, C. E. 1956. Yerba de la fleche -- arrow and fish poison of the American Southwest. *Econ. Bot.* 10: 362-366.

Bryn, T. K. 1963. Curare: its history and usage. J. B. Lippincott. Philadelphia, PA.

Burnap, T. K. & D. M. Little, Jr. (Editors). 1968. The flying death. Classical papers and commentary on curare. Inter. Anesthesiology Clinics. Little, Brown, and Co. Boston, MA. 6(2): 399-739.

Cheney, R. H. 1926. The ancient and modern use of plant arrow poisons. *Sci. Monthly* 23: 552-555.

Cheney, R. H. 1926. Plant arrow poisons: their sources, preparation, and effects. *J. New York Bot.*

Gard. 27: 174-177.

Cheney, R. H. 1931. Geographic and taxonomic distribution of American plant arrow poisons. *American J. Bot.* 18: 136-145.

Kopp, B. et al. 1992. Analysis of some Malaysian dart poisons. *J. Ethnopharm.* 36(1): 57-62.

Krukoff, B. A. 1937. Notes on the botanical components of curare. *Bull. Torrey Bot. Club* 64: 401-409.

Krukoff, B. A. & H. N. Moldenke. 1938. Studies of American Menispermaceae, with special reference to species used in the preparation of arrow poisons. *Brittonia* 3: 1-74 + suppl. 1-5.

Maitai, C. K. et al. 1973. A survey on the use of poisoned arrows in Kenya during the period 1964-1971. *East African Med. J.* 50: 100-104.

Reuck, A. V. S. de (editor). 1962. Curare and curare-like agents. Ciba Found. Study Group No. 12. Little, Brown and Co. Boston, MA. 103 pp.

Thomas, K. B. 1963. Curare: its history and usage. J. B. Lippincott Co. Philadelphia, PA. 144 pp.

Vellard, J. 1965. Historie du curare. Gallimard. Paris, France.

FISH POISONS

Acevedo-Rodriguez, P. 1990. The occurrence of piscicides and stupefactans in the plant kingdom. *Adv. Econ. Bot.* 8: 1-23.

Bradley, C. E. 1956. Yerba de la fleche -- arrow and fish poisons of the American Southwest. *Econ. Bot.* 10: 362-366.

Brandt, A. von. 1972. Fish catching methods of the world. Fishing News (Books), Ltd. London, England. Pp. 22-25.

Heizer, R. F. 1941. The use of plants for fish poisoning by the California Indians. *Leaflets. Western Bot.* 3(2): 43, 44.

Heizer, R. F. 1949. Fish poisons. In, Stewart, J. H. (Editor). *Handbook of South American Indians. The*

comparative ethnology of South American Indians. Smithsonian Institution Bureau of American Ethnology. Bulletin 143. 5: 565-586.

Heizer, R. F. 1953. Aboriginal fish poisons. Bureau of American Ethnology. Bull. 151. Anthropological Papers, No. 38. Pp. 225-283.

Higbee, E. C. 1947. *Lonchocarpus* -- a fish poison insecticide. Econ. Bot. 1: 427-436.

Howes, F. N. 1930. Fish-poison plants. Kew Bull. Misc. Inform. 4: 129-153.

Kamen-Kaye, D. 1977. Ichthyotoxic plants and the term "barbasco." Bot. Mus. Leaflets. Harvard Univ. 25: 71-90.

Killip, E. P. & A. C. Smith. 1931. The use of fish poisons in South America. Ann. Rep. Smithsonian Inst., 1930. Washington, D. C. Pp. 401-408.

McFarland, J. W. 1951. Poisonous plants used for fishing. Yosemite Nature Notes 30: 14-21.

Neuwinger, H. D. 1994. Fish poisoning plants in Africa. Bot. Acta 107: 264-270.

Nishimoto, S. K. 1969. Plants used as fish poisons. Newsletter Hawaiian Bot. Soc. 8: 20-23.

Pal, D. C. & A. M. Saren. 1986. Some plants used by tribes of India for poisoning/stupefying fishes. J. Econ. Tax. Bot. 8: 13-16.

Pennington, C. W. 1957. Tarahumar fish stupefaction plants. Econ. Bot. 12(1): 95-102.

Quigley, C. 1956. Aboriginal fish poisons and the diffusion problem. American Anthrop. 58: 508-525.

Rickard, P. & P. A. Cox. 1986. Use of *Derris* as a fish poison in the Solomon Islands. Econ. Bot. 40: 479-484.

Stokes, J. F. G. 1921. Fish-poisoning in the Hawaiian Islands with notes on the custom in southern Polynesia. Occas. Pap. Bernice P. Bishop Mus. 7(10): 219-236.

Wilhelm, G. 1974. The mullein: plant piscicide of the mountain folk culture. Geogr. Rev. April: 235-252.

INSECT POISONS

Annason, J. T. et al. 1989. Insecticides of plant origin. American Chem. Soc. Washington, D. C. 213 pp.

Barnes, D. K. & R. H. Freyre. 1966. Recovery of natural insecticides from *Tephrosia vogelii*. II. Toxicological properties of rotenoids extracted from fresh and oven-dried leaves. Econ. Bot. 20(4): 368-371.

Casida, J. E. & G. B. Quistad. 1995. Pyrethrum flowers: production, chemistry, toxicology, and uses. Oxford Univ. Press. New York, NY. 350 pp.

Crosby, B. G. 1971. Minor insecticides of plant origin. In, Jacobson, M. & D. G. Crosby. Naturally occurring insecticides. Marcel Dekker. New York, NY. Pp. 177-239.

Duffey, S. S. 1980. Sequestration of plant natural-products by insects. Ann. Rev. Entom. 25: 447-477.

Grainge, M. & S. Ahmed. 1988. Handbook of plants with pest-control properties. John Wiley & Sons. New York, NY.

Higbee, E. C. 1947. *Lonchocarpus* -- a fish poison insecticide. Econ. Bot. 1: 427-436.

Isman, M. B. 1994. Botanical insecticides. Pesticide Outlook 5: 26-31.

Jacobson, M. 1958. Insecticides from plants. A review of the literature, 1941-1953. Agric. Handbook No. 154. U. S. Dept. Agric. Washington, D. C. 299 pp.

Jacobson, M. 1975. Insecticides from plants. A review of the literature, 1954-1971. U. S. Dept. Agric. Washington, D. C. 138 pp.

Jacobson, M. & D. G. Crosby (editors). 1971. Naturally occurring insecticides. Marcel Dekker. New York, NY.

Klocke, J. A. 1989. Plant compounds as sources and models of insect-control agents. Econ. Med. Plant Res. 3: 104-144.

Lydon, J. & S. O. Duke. 1989. The potential of pesticides from plants. In, Craker & Simon, Pp. 1-41.

Metcalf, R. L. 1977. Plant derivatives for insect control. In, Seigler, D. S. (Editor). Crop resources. Academic Press. New York, NY. Pp. 165-177.

Nathanson, J. A. 1984. Caffeine and related methylxanthines: possible naturally occurring pesticides. Science 226: 184-187.

Run, P. J. van. 1974. The production of pyrethrum. Trop. Abstr. 29: 237-244.

ORDEAL POISONS

Bisset, N. G. & J. M. Leeuwenberg. 1968. The use of *Strychnos* in Central African ordeal and arrow poisons. Lloydia 31: 208.

Holmstedt, B. 1972. The ordeal bean of Old Calabar. In, Swain, T. (editor). Plants in the development of modern medicine. Harvard Univ. Press. Cambridge, MA. 303-360.

Marwick, M. G. 1963. A note on ordeal poisoning in east central Africa. Man 63: 45, 46.

Robb, G. L. 1957. The ordeal poisons of Madagascar and Africa. Bot. Mus. Leaflets. Harvard Univ. 17: 265-316.